

AN EXTENSION THEOREM FOR SEPARATELY HOLOMORPHIC FUNCTIONS WITH SINGULARITIES

MAREK JARNICKI (KRAKÓW)^(†) PETER PFLUG (OLDENBURG)^(‡)

ABSTRACT. Let $D_j \subset \mathbb{C}^{k_j}$ be a pseudoconvex domain and let $A_j \subset D_j$ be a locally pluripolar set, $j = 1, \dots, N$. Put

$$X := \bigcup_{j=1}^N A_1 \times \dots \times A_{j-1} \times D_j \times A_{j+1} \times \dots \times A_N \subset \mathbb{C}^{k_1 + \dots + k_N}.$$

Let U be an open connected neighborhood of X and let $M \subsetneq U$ be an analytic subset. Then there exists an analytic subset $\widehat{M}$ of the ‘envelope of holomorphy’ $\widehat{X}$ of X with $\widehat{M} \cap X \subset M$ such that for every function f separately holomorphic on $X \setminus M$ there exists an $\widehat{f}$ holomorphic on $\widehat{X} \setminus \widehat{M}$ with $\widehat{f}|_{X \setminus M} = f$. The result generalizes special cases which were studied in [Ökt 1998], [Ökt 1999], [Sic 2000], and [Jar-Pfl 2001].

1. Introduction. Main Theorem. Let $N \in \mathbb{N}$, $N \geq 2$, and let

$$\emptyset \neq A_j \subset D_j \subset \mathbb{C}^{k_j},$$

where D_j is a domain, $j = 1, \dots, N$. We define an N -fold cross

$$\begin{aligned} X &:= \mathbb{X}(A_1, \dots, A_N; D_1, \dots, D_N) \\ &:= \bigcup_{j=1}^N A_1 \times \dots \times A_{j-1} \times D_j \times A_{j+1} \times \dots \times A_N \subset \mathbb{C}^{k_1 + \dots + k_N}. \end{aligned} \quad (1)$$

Observe that X is connected.

Let $\Omega \subset \mathbb{C}^n$ be an open set and let $A \subset \Omega$. Put

$$h_{A, \Omega} := \sup\{u : u \in \mathcal{PSH}(\Omega), u \leq 1 \text{ on } \Omega, u \leq 0 \text{ on } A\},$$

1991 *Mathematics Subject Classification.* 32D15, 32D10.

^(†) Research partially supported by the KBN grant No. 5 P03A 033 21.

^(‡) Research partially supported by the Niedersächsisches Ministerium für Wissenschaft und Kultur, Az. 15.3 – 50 113(55) PL.

where $\mathcal{PSH}(\Omega)$ denotes the set of all functions plurisubharmonic on Ω . Define

$$\omega_{A,\Omega} := \lim_{k \rightarrow +\infty} h_{A \cap \Omega_k, \Omega_k}^*,$$

where $(\Omega_k)_{k=1}^\infty$ is a sequence of relatively compact open sets $\Omega_k \subset \Omega_{k+1} \subset \subset \Omega$ with $\bigcup_{k=1}^\infty \Omega_k = \Omega$ (h^* denotes the upper semicontinuous regularization of h). Observe that the definition is independent of the chosen exhausting sequence $(\Omega_k)_{k=1}^\infty$. Moreover, $\omega_{A,\Omega} \in \mathcal{PSH}(\Omega)$.

For an N -fold cross $X = \mathbb{X}(A_1, \dots, A_N; D_1, \dots, D_N)$ put

$$\widehat{X} := \{(z_1, \dots, z_N) \in D_1 \times \dots \times D_N : \sum_{j=1}^N \omega_{A_j, D_j}(z_j) < 1\};$$

notice that $\widehat{X}$ may be empty. Observe that $\widehat{X}$ is pseudoconvex if $D_1, \dots, D_N$ are pseudoconvex domains.

We say that a subset $\emptyset \neq A \subset \mathbb{C}^n$ is *locally pluriregular* if $h_{A \cap \Omega, \Omega}^*(a) = 0$ for any $a \in A$ and for any open neighborhood Ω of a (in particular, $A \cap \Omega$ is non-pluripolar).

Note that if $A_1, \dots, A_N$ are locally pluriregular, then $X \subset \widehat{X}$ and, moreover, $\widehat{X}$ is connected (Lemma 5).

Let U be a connected neighborhood of X and let $M \subsetneq U$ be an analytic subset (M may be empty). We say that a function $f: X \setminus M \rightarrow \mathbb{C}$ is *separately holomorphic* ($f \in \mathcal{O}_s(X \setminus M)$) if for any $(a_1, \dots, a_N) \in A_1 \times \dots \times A_N$ and $j \in \{1, \dots, N\}$ the function $f(a_1, \dots, a_{j-1}, \cdot, a_{j+1}, \dots, a_N)$ is holomorphic in the domain $\{z_j \in D_j : (a_1, \dots, a_{j-1}, z_j, a_{j+1}, \dots, a_N) \notin M\}$.

The main result of our paper is the following extension theorem for separately holomorphic functions.^(†)

Main Theorem. *Let $D_j \subset \mathbb{C}^{k_j}$ be a pseudoconvex domain and let $A_j \subset D_j$ be a locally pluriregular set, $j = 1, \dots, N$. Let $M \subsetneq U$ be an analytic subset of an open connected neighborhood U of $X = \mathbb{X}(A_1, \dots, A_N; D_1, \dots, D_N)$ (M may be empty). Then there exists a pure one-codimensional analytic subset $\widehat{M} \subset \widehat{X}$ such that:*

- $\widehat{M} \cap U_0 \subset M$ for an open neighborhood U_0 of X , $U_0 \subset U$,
- for every $f \in \mathcal{O}_s(X \setminus M)$ there exists exactly one $\widehat{f} \in \mathcal{O}(\widehat{X} \setminus \widehat{M})$ with $\widehat{f}|_{X \setminus M} = f$.

Moreover, if $U = \widehat{X}$, then we can take $\widehat{M} :=$ the union of all one-codimensional irreducible components of M .

The proof will be given in Sections 3 (the case $U = \widehat{X}$) and 4 (the general case).

^(†) We like to thank Professor Józef Siciak for turning our attention to the problem.

Remark. Notice that the Main Theorem may be generalized to the case where D_j is Riemann–Stein domains over $\mathbb{C}^{k_j}$, $j = 1, \dots, N$.

Observe that in the case $M = \emptyset$, $N = 2$, the Main Theorem is nothing else than the following cross theorem.

Theorem 1 (cf. [Ale-Zer 2001]). *Let $D \subset \mathbb{C}^p$, $G \subset \mathbb{C}^q$ be pseudoconvex domains and let $A \subset D$, $B \subset G$ be locally pluriregular. Put $X := \mathbb{X}(A, B; D, G)$. Then for any $f \in \mathcal{O}_s(X)$ there exists exactly one $\hat{f} \in \mathcal{O}(\hat{X})$ with $\hat{f} = f$ on X .*

Remark. (a) $M = \emptyset$: There is a long list of papers discussing the case $N = 2$ (under various assumptions): [Sic 1969], [Zah 1976], [Sic 1981], [Shi 1989], [Ngu-Zer 1991], [Ngu 1997], [Ale-Zer 2001]. The case $N \geq 2$, $k_1 = \dots = k_N = 1$ can be found in [Sic 1981]. The general case $N \geq 2$, $k_1, \dots, k_N \geq 1$ was solved in [Ngu-Zer 1995] ^(†).

(b) $M \neq \emptyset$: J. Siciak [Sic 2000] solved the case: $N \geq 2$, $k_1 = \dots = k_N = 1$, $D_1 = \dots = D_N = \mathbb{C}$, $M = P^{-1}(0)$, where P is a non-zero polynomial of N complex variables. The special subcase $N = 2$, $P(z, w) := z - w$ had been studied in [Ökt 1998]. The general case for $N = 2$, $k_1 = k_2 = 1$ was solved in [Jar-Pfl 2001]; see also [Ökt 1999] for a partial discussion of the case $N = 2$, $k_1, k_2 \geq 1$.

It remains an open problem what happens if the singular set M is assumed to be, for instance, a pluripolar relatively closed subset of U (see also [Chi-Sad 1988]). The following particular case will be proved in Section 5.

Theorem 2. *Let $A, B \subset \mathbb{C}$ be locally regular. Let $M \subset \mathbb{C}^2$ be a closed pluripolar set and let $X := \mathbb{X}(A, B; \mathbb{C}, \mathbb{C})$. Then there exist a closed pluripolar set $\widehat{M} \subset \mathbb{C}^2$ and $A' \subset A$, $B' \subset B$ with $A \setminus A'$, $B \setminus B'$ being polar, such that for every $f \in \mathcal{O}_s(X \setminus M)$ there exists exactly one $\hat{f} \in \mathcal{O}(\mathbb{C}^2 \setminus \widehat{M})$ with $\hat{f}|_{X' \setminus (M \cup \widehat{M})} = f|_{X' \setminus (M \cup \widehat{M})}$, where $X' := \mathbb{X}(A', B'; \mathbb{C}, \mathbb{C})$.*

Notice that Theorem 2 may be thought as a partial generalization of [Sic 2000].

2. Auxiliary results. The following lemma gathers a few standard results, which will be frequently used in the sequel.

Lemma 3 (cf. [Kli 1991], [Jar-Pfl 2000], § 3.5). (a) *Let $\Omega \subset \mathbb{C}^n$ be a bounded open set and let $A \subset \Omega$. Then:*

- *If $P \subset \mathbb{C}^n$ is pluripolar, then $h_{A \setminus P, \Omega}^* = h_{A, \Omega}^*$.*
- *$h_{A_k \cap \Omega_k, \Omega_k}^* \searrow h_{A, \Omega}^*$ (pointwise on Ω) for any sequence of open sets $\Omega_k \nearrow \Omega$ and any sequence $A_k \nearrow A$.*
- *$\omega_{A, \Omega} = h_{A, \Omega}^*$.*
- *The following conditions are equivalent:*
for any connected component S of Ω the set $A \cap S$ is non-pluripolar;
 $h_{A, \Omega}^*(z) < 1$ for any $z \in \Omega$.

^(†) We like to thank Professor Nguyen Thanh Van for calling our attention to that paper.

• If A is non-pluripolar, $0 < \alpha < 1$, and $\Omega_\alpha := \{z \in \Omega : h_{A,\Omega}^*(z) < \alpha\}$, then for any connected component S of Ω_α the set $A \cap S$ is non-pluripolar (in particular, $A \cap S \neq \emptyset$).

• If A is locally pluriregular, $0 < \alpha \leq 1$, and $\Omega_\alpha := \{z \in \Omega : h_{A,\Omega}^*(z) < \alpha\}$, then $h_{A,\Omega}^* = \alpha h_{A,\Omega_\alpha}^*$ on Ω_α .

(b) Let $\Omega \subset \mathbb{C}^n$ be an open set and let $A \subset \Omega$. Then:

- $\omega_{A,\Omega} \in \mathcal{PSH}(\Omega)$.
- If A is locally pluriregular, then $\omega_{A,\Omega}(a) = 0$ for any $a \in A$.
- If $P \subset \mathbb{C}^n$ is pluripolar, then $\omega_{A \setminus P, \Omega} = \omega_{A,\Omega}$.
- If A is locally pluriregular and $P \subset \mathbb{C}^n$ is pluripolar, then $A \setminus P$ is locally pluriregular.

Moreover, we get:

Lemma 4. (a) Let $A_j \subset \mathbb{C}^{k_j}$ be locally pluriregular, $j = 1, \dots, N$, then $A_1 \times \dots \times A_N$ is locally pluriregular.

(b) Let $A_j \subset \Omega_j \Subset \mathbb{C}^{k_j}$, Ω_j a domain, A_j locally pluriregular, $j = 1, \dots, N$, $N \geq 2$. Put

$$\Omega := \{(z_1, \dots, z_N) \in \Omega_1 \times \dots \times \Omega_N : \sum_{j=1}^N h_{A_j, \Omega_j}^*(z_j) < 1\}$$

(observe that $A_1 \times \dots \times A_N \subset \Omega$). Then

$$h_{A_1 \times \dots \times A_N, \Omega}^* = \sum_{j=1}^N h_{A_j, \Omega_j}^* \quad \text{on } \Omega.$$

Proof. (a) is an immediate consequence of the product property for the relatively extremal function

$$h_{B_1 \times \dots \times B_N, \Omega_1 \times \dots \times \Omega_N}^* = \max\{h_{B_j, \Omega_j}^* : j = 1, \dots, N\};$$

cf. [Ngu-Sic 1991].

(b) First observe that

$$\sum_{j=1}^N h_{A_j, \Omega_j}^* \leq h_{A_1 \times \dots \times A_N, \Omega}^* \quad \text{on } \Omega.$$

To get the opposite inequality we proceed by induction on $N \geq 2$.

Let $N = 2$: The proof of this step is taken from [Sic 1981]. For the reader's convenience we repeat the details.

Put $u := h_{A_1 \times A_2, \Omega}^* \in \mathcal{PSH}(\Omega)$ and fix a point $(a_1, a_2) \in \Omega$.

If $a_1 \in A_1$ (thus $h_{A_1, \Omega_1}^*(a_1) = 0$), then $u(a_1, \cdot) \in \mathcal{PSH}(\Omega_2)$ with $u(a_1, \cdot) \leq 1$ and $u(a_1, \cdot) \leq 0$ on A_2 . Therefore,

$$u(a_1, \cdot) \leq h_{A_2, \Omega_2}^* = h_{A_1, \Omega_1}^*(a_1) + h_{A_2, \Omega_2}^* \quad \text{on } \Omega_2.$$

In particular, $u(a_1, a_2) \leq h_{A_1, \Omega_1}^*(a_1) + h_{A_2, \Omega_2}^*(a_2)$.

Observe that the same argument shows that if $a_2 \in A_2$, then $u(\cdot, a_2) \leq h_{A_1, \Omega_1}^*$ on Ω_1 .

If $a_1 \notin A_1$, then $h_{A_1, \Omega_1}^*(a_1) + h_{A_2, \Omega_2}^*(a_2) < 1$ and therefore $\alpha := 1 - h_{A_2, \Omega_2}^*(a_2) \in (0, 1]$. Put

$$(\Omega_2)_\alpha := \{z_2 \in \Omega_2 : h_{A_2, \Omega_2}^*(z_2) < \alpha\}.$$

It is clear that $A_2 \subset (\Omega_2)_\alpha \ni a_2$. Put

$$v := \frac{1}{\alpha} \left(u(a_1, \cdot) - h_{A_1, \Omega_1}^*(a_1) \right) \in \mathcal{PSH}((\Omega_2)_\alpha).$$

Then $v \leq 1$ and $v \leq 0$ on A_2 . Therefore, by Lemma 3(a),

$$v \leq h_{A_2, (\Omega_2)_\alpha}^*(a_2) = \frac{1}{\alpha} h_{A_2, \Omega_2}^*(a_2) \quad \text{on } (\Omega_2)_\alpha.$$

Consequently, $u(a_1, a_2) \leq h_{A_1, \Omega_1}^*(a_1) + h_{A_2, \Omega_2}^*(a_2)$, which finishes the proof for $N = 2$.

Now, assume that the formula is true for $N - 1 \geq 2$. Put

$$\tilde{\Omega} := \{(z_1, \dots, z_{N-1}) \in \Omega_1 \times \dots \times \Omega_{N-1} : \sum_{j=1}^{N-1} h_{A_j, \Omega_j}^*(z_j) < 1\}$$

and fix an arbitrary $z = (\tilde{z}, z_N) \in \Omega$. Obviously, $\tilde{z} \in \tilde{\Omega}$. In virtue of the inductive hypothesis, we conclude that

$$h_{A_1 \times \dots \times A_{N-1}, \tilde{\Omega}}^*(\tilde{z}) = \sum_{j=1}^{N-1} h_{A_j, \Omega_j}^*(z_j). \quad (2)$$

Now we apply the case $N = 2$ to the following situation:

$$\Omega' := \{(\tilde{w}, w_N) \in \tilde{\Omega} \times \Omega_N : h_{A_1 \times \dots \times A_{N-1}, \tilde{\Omega}}^*(\tilde{w}) + h_{A_N, \Omega_N}^*(w_N) < 1\}.$$

So

$$h_{A_1 \times \dots \times A_{N-1}, \tilde{\Omega}}^*(\tilde{w}) + h_{A_N, \Omega_N}^*(w_N) = h_{A_1 \times \dots \times A_N, \Omega'}^*(\tilde{w}, w_N), \quad (\tilde{w}, w_N) \in \Omega'.$$

Note that $\Omega' = \Omega$. Hence

$$h_{A_1 \times \dots \times A_N, \Omega}^*(\tilde{z}, z_N) = h_{A_1 \times \dots \times A_{N-1}, \tilde{\Omega}}^*(\tilde{z}) + h_{A_N, \Omega_N}^*(z_N) \stackrel{(2)}{=} \sum_{j=1}^N h_{A_j, \Omega_j}^*(z_j).$$

□

Lemma 5. *Let $X = \mathbb{X}(A_1, \dots, A_N; D_1, \dots, D_N)$ be an N -fold cross as in (1). If $A_1, \dots, A_N$ are locally pluriregular, then $\widehat{X}$ is a domain.*

Proof. Using exhaustion by bounded domains we may assume that the D_j 's are bounded.

We know that $X \subset \widehat{X}$. Let $z^0 = (z_1^0, \dots, z_N^0) \in \widehat{X}$ be an arbitrary point.

If $\sum_{j=2}^N h_{A_j, \Omega_j}^*(z_j^0) = 0$, then $D_1 \times \{(z_2^0, \dots, z_N^0)\} \subset \widehat{X}$. Therefore, z^0 can be joined inside $D_1 \times \{(z_2^0, \dots, z_N^0)\}$ with $(a_1, z_2^0, \dots, z_N^0)$ for some $a_1 \in A_1$.

If $\sum_{j=2}^N h_{A_j, \Omega_j}^*(z_j^0) =: \varepsilon > 0$, put

$$(D_1)_{1-\varepsilon} := \{z_1 \in D_1 : h_{A_1, D_1}^*(z_1) < 1 - \varepsilon\}.$$

Then, in virtue of Lemma 3(a), the connected component S of $(D_1)_{1-\varepsilon}$, that contains z_1^0 , intersects A_1 . Therefore, z^0 can be joined inside $S \times \{(z_2^0, \dots, z_N^0)\} \subset \widehat{X}$ with $(a_1, z_2^0, \dots, z_N^0)$ for some $a_1 \in A_1$.

Now we repeat the above argument for the second component of the point $(a_1, z_2^0, \dots, z_N^0)$. Finally, the point z^0 can be joined inside $\widehat{X}$ with $(a_1, \dots, a_N) \in A_1 \times \dots \times A_N \subset X$. Since X is connected, the proof is completed. $\square$

Lemma 6 (Identity theorems). (a) *Let $\Omega \subset \mathbb{C}^n$ be a domain and let $A \subset \Omega$ be non-pluripolar. Then any $f \in \mathcal{O}(\Omega)$ with $f|_A = 0$ vanishes identically on Ω .*

(b) *Let $D \subset \mathbb{C}^p, G \subset \mathbb{C}^q$ be domains, let $A \subset D, B \subset G$ be locally pluriregular sets, and let $X := \mathbb{X}(A, B; D, G)$. Let $M \subsetneq U$ be an analytic subset of an open connected neighborhood U of X . Assume that $A' \subset A, B' \subset B$ are such that:*

- $A \setminus A'$ and $B \setminus B'$ are pluripolar (in particular, A', B' are also locally pluriregular),
- $M_z := \{w \in G : (z, w) \in M\} \neq G$ for any $z \in A'$,
- $M^w := \{z \in D : (z, w) \in M\} \neq D$ for any $w \in B'$.

Then:

- (b₁) *If $f \in \mathcal{O}_s(X \setminus M)$ and $f = 0$ on $A' \times B' \setminus M$ ^(†), then $f = 0$ on $X \setminus M$.*
- (b₂) *If $g \in \mathcal{O}(U \setminus M)$ and $g = 0$ on $A' \times B' \setminus M$, then $g = 0$ on $U \setminus M$.*

Proof. (a) is obvious.

(b₁) Take a point $(a_0, b_0) \in X \setminus M$. We may assume that $a_0 \in A$. Since $A \setminus A'$ is pluripolar, there exists a sequence $(a_k)_{k=1}^\infty \subset A'$ such that $a_k \rightarrow a_0$. The set $Q := \bigcup_{k=0}^\infty M_{a_k}$ is pluripolar. Consequently, the set $B'' := B' \setminus Q$ is non-pluripolar. We have $f(a_k, w) = 0, w \in B'', k = 1, 2, \dots$. Hence $f(a_0, w) = 0$ for any $w \in B''$. Finally, $f(a_0, w) = 0$ on $G \setminus M_{a_0} \ni b_0$.

(b₂) Take an $a_0 \in A'$. Since $M_{a_0} \neq G$, there exists a $b_0 \in B' \setminus M_{a_0}$. Let $P = \Delta_{a_0}(r) \times \Delta_{b_0}(r) \subset U \setminus M$ ($\Delta_{z_0}(r) \subset \mathbb{C}^p$ denotes the polydisc with center $z_0 \in \mathbb{C}^p$ and radius $r > 0$). Then $g(\cdot, w) = 0$ on $A' \cap \Delta_{a_0}(r)$ for any $w \in B' \cap \Delta_{b_0}(r)$. The set $A' \cap \Delta_{a_0}(r)$ is non-pluripolar. Hence $g(\cdot, w) = 0$ on $\Delta_{a_0}(r)$ for any $w \in B' \cap \Delta_{b_0}(r)$.

^(†) Here and in the sequel to simplify notation we write $P \times Q \setminus M$ instead of $(P \times Q) \setminus M$.

By the same argument for the second variable we get $g = 0$ on P and, consequently, on U . $\square$

3. Proof of the Main Theorem in the case where $U = \widehat{X}$. We proceed by several reduction steps. First observe that, by Lemma 6(a), the function $\widehat{f}$ is uniquely determined (if exists).

Step 1. *To prove the Main Theorem for $M \neq \emptyset$ it suffices to consider only the case where M is pure one-codimensional.*

Proof of Step 1. Since $\widehat{X}$ is pseudoconvex, the arbitrary analytic set $M \subset \widehat{X}$ can be written as

$$M = \{z \in \widehat{X} : g_1(z) = \cdots = g_k(z) = 0\},$$

where $g_j \in \mathcal{O}(\widehat{X})$, $g_j \not\equiv 0$, $j = 1, \dots, k$. Then $M_j := g_j^{-1}(0)$ is pure one-codimensional.

Take an $f \in \mathcal{O}_s(X \setminus M)$. Observe that $f_j := f|_{X \setminus M_j} \in \mathcal{O}_s(X \setminus M_j)$. By the reduction assumption there exists an $\widehat{f}_j \in \mathcal{O}(\widehat{X} \setminus M_j)$ such that $\widehat{f}_j = f$ on $X \setminus M_j$. In virtue of Lemma 6(a), gluing the functions $(\widehat{f}_j)_{j=1}^k$, leads to an $\widehat{f} \in \mathcal{O}(\widehat{X} \setminus M)$ with $\widehat{f} = \widehat{f}_j$ on $\widehat{X} \setminus M_j$, $j = 1, \dots, k$. Therefore, $\widehat{f} = f$ on $X \setminus M$.

Finally, since $\text{codim}(M \setminus \widehat{M}) \geq 2$, the function $\widehat{f}$ extends holomorphically across $M \setminus \widehat{M}$. $\square$

From now on we assume that M is empty or pure one-codimensional.

Step 2. *To prove the Main Theorem it suffices to consider only the case where M is empty or pure one-codimensional and $D_1, \dots, D_N$ are bounded pseudoconvex.*

Proof of Step 2. Let $D_1, \dots, D_N$ be arbitrary pseudoconvex domains. Let $D_{j,k} \nearrow D_j$, $D_{j,k} \Subset D_j$, where $D_{j,k}$ are pseudoconvex domains with $A_{j,k} := A_j \cap D_{j,k} \neq \emptyset$. Observe that all the $A_{j,k}$'s are locally pluriregular. Put

$$X_k := \mathbb{X}(A_{1,k}, \dots, A_{N,k}; D_{1,k}, \dots, D_{N,k}) \subset X;$$

note that $\widehat{X}_k \nearrow \widehat{X}$.

Let $f \in \mathcal{O}_s(X \setminus M)$ be given. By the reduction assumption, for each k there exists an $\widehat{f}_k \in \mathcal{O}(\widehat{X}_k \setminus M)$ with $\widehat{f}_k = f$ on $X_k \setminus M$. By Lemma 6(a), $\widehat{f}_{k+1} = \widehat{f}_k$ on $\widehat{X}_k \setminus M$. Therefore, gluing the $\widehat{f}_k$'s, we obtain an $\widehat{f} \in \mathcal{O}(\widehat{X} \setminus M)$ with $\widehat{f} = f$ on $X \setminus M$. $\square$

From now on we assume that M is empty or pure one-codimensional and $D_1, \dots, D_N$ are bounded pseudoconvex.

Step 3. *To prove the Main Theorem it suffices to consider only the case $N = 2$.*

Remark. In virtue of Theorem 1, Step 3 finishes the proof of the Main Theorem for $M = \emptyset$.

Proof of Step 3. We proceed by induction on $N \geq 2$. Suppose that the Main Theorem is true for $N - 1 \geq 2$. We have to discuss the case of an N -fold cross $X = \mathbb{X}(A_1, \dots, A_N; D_1, \dots, D_N)$, where $D_1, \dots, D_N$ are bounded pseudoconvex. Let $M \subset \widehat{X}$ be empty or pure one-codimensional.

Let $f \in \mathcal{O}_s(X \setminus M)$.

Observe that

$$X = (Y \times A_N) \cup (\widehat{A} \times D_N),$$

where

$$Y := \mathbb{X}(A_1, \dots, A_{N-1}; D_1, \dots, D_{N-1}), \quad \widehat{A} := A_1 \times \dots \times A_{N-1}.$$

We also mention that for any $a_N \in A_N$ we have

$$\{(z_1, \dots, z_{N-1}) \in \mathbb{C}^{k_1} \times \dots \times \mathbb{C}^{k_{N-1}} : (z_1, \dots, z_{N-1}, a_N) \in \widehat{X}\} = \widehat{Y}.$$

Now fix an $a_N \in A_N$ such that

$$M_{a_N} := \{(z_1, \dots, z_{N-1}) \in \widehat{Y} : (z_1, \dots, z_{N-1}, a_N) \in M\} \subsetneq \widehat{Y};$$

in particular, M_{a_N} is empty or one-codimensional (in $\widehat{Y}$). Recall that $A_1, \dots, A_{N-1}$ are locally pluriregular. By inductive assumption there exists an $\widehat{f}_{a_N} \in \mathcal{O}(\widehat{Y} \setminus M_{a_N})$ with $\widehat{f}_{a_N} = f(\cdot, a_N)$ on $Y \setminus M_{a_N}$.

To continue define the following 2-fold cross

$$Z := \mathbb{X}(\widehat{A}, A_N; \widehat{Y}, D_N).$$

Notice that Z satisfies all the properties for the case $N = 2$: $\widehat{Y}, D_N$ are bounded pseudoconvex domains, $\widehat{A} \subset \widehat{Y}$, $A_N \subset D_N$ are locally pluriregular.

In virtue of Lemma 4, we have

$$\widehat{Z} = \{(\widehat{z}, z_N) \in \widehat{Y} \times D_N : h_{\widehat{A}, \widehat{Y}}^*(\widehat{z}) + h_{A_N, D_N}^*(z_N) < 1\} = \widehat{X}.$$

Define $\widetilde{f}: Z \setminus M \rightarrow \mathbb{C}$,

$$\widetilde{f}(z) = \widetilde{f}(\widehat{z}, z_N) := \begin{cases} \widehat{f}_{z_N}(\widehat{z}) & \text{if } z \in \widehat{Y} \times A_N \\ f(z) & \text{if } z \in \widehat{A} \times D_N \end{cases}.$$

Obviously, $\widetilde{f}$ is well-defined and therefore $\widetilde{f} \in \mathcal{O}_s(Z \setminus M)$.

Using the case $N = 2$, we find another function $\widehat{f} \in \mathcal{O}(\widehat{Z} \setminus M)$ with $\widehat{f} = \widetilde{f}$ on $Z \setminus M$. Recall that $\widehat{Z} = \widehat{X}$. Hence $\widehat{f} = f$ on $X \setminus M$. $\square$

What remains is to prove the case $N = 2$ and $M \neq \emptyset$. From now on we simplify our notation and we are looking for the following configuration:

Let $A \subset D \Subset \mathbb{C}^p$, $B \subset G \Subset \mathbb{C}^q$, where D, G are bounded pseudoconvex domains, A, B are locally pluriregular. Put, as always,

$$X := \mathbb{X}(A, B; D, G), \quad \widehat{X} := \{(z, w) \in D \times G : h_{A, D}^*(z) + h_{B, G}^*(w) < 1\}.$$

Moreover, let M be a pure one-codimensional analytic subset of $\widehat{X}$.

We like to show that any $f \in \mathcal{O}_s(X \setminus M)$ extends holomorphically to $\widehat{X} \setminus M$.

Step 4. Let X , M , and f be as above. Let $(D_j)_{j=1}^\infty$, $(G_j)_{j=1}^\infty$ be sequences of pseudoconvex domains, $D_j \Subset D$, $G_j \Subset G$, with $D_j \nearrow D$, $G_j \nearrow G$. Moreover, let $A' \subset A$, $B' \subset B$ be such that $A \setminus A'$, $B \setminus B'$ are pluripolar, and $A' \cap D_j \neq \emptyset$, $B' \cap G_j \neq \emptyset$, $j \in \mathbb{N}$. For each $j \in \mathbb{N}$ we assume that:

for any $(a, b) \in (A' \cap D_j) \times (B' \cap G_j)$ there exist polydiscs $\Delta_a(r_{a,j}) \subset D_j$, $\Delta_b(s_{b,j}) \subset G_j$ with $(\Delta_a(r_{a,j}) \times G_j) \cup (D_j \times \Delta_b(s_{b,j})) \subset \widehat{X}$, and functions $f_{a,j} \in \mathcal{O}(\Delta_a(r_{a,j}) \times G_j \setminus M)$, $f^{b,j} \in \mathcal{O}(D_j \times \Delta_b(s_{b,j}) \setminus M)$ such that

- $f_{a,j} = f$ on $(A' \cap \Delta_a(r_{a,j})) \times G_j \setminus M$,
- $f^{b,j} = f$ on $D_j \times (B' \cap \Delta_b(s_{b,j})) \setminus M$.

Then there exists an $\widehat{f} \in \mathcal{O}(\widehat{X} \setminus M)$ with $\widehat{f} = f$ on $X \setminus M$.

Proof of Step 4. Fix a $j \in \mathbb{N}$. Put

$$\begin{aligned} \widetilde{U}_j &:= \bigcup_{\substack{a \in A' \cap D_j \\ b \in B' \cap G_j}} (\Delta_a(r_{a,j}) \times G_j) \cup (D_j \times \Delta_b(s_{b,j})), \\ X'_j &:= ((A \cap D_j) \times G_j) \cup (D_j \times (B \cap G_j)). \end{aligned}$$

Note that

$$X'_j := ((A' \cap D_j) \times G_j) \cup (D_j \times (B' \cap G_j)) \subset \widetilde{U}_j.$$

We like to glue the functions $(f_{a,j})_{a \in A' \cap D_j}$ and $(f^{b,j})_{b \in B' \cap G_j}$ to obtain a global holomorphic function f_j on $\widetilde{U}_j \setminus M$:

Let $a \in A' \cap D_j$, $b \in B' \cap G_j$. Observe that

$$\begin{aligned} f_{a,j} &= f \quad \text{on } (A' \cap \Delta_a(r_{a,j})) \times G_j \setminus M, \\ f^{b,j} &= f \quad \text{on } D_j \times (B' \cap \Delta_b(s_{b,j})) \setminus M. \end{aligned}$$

Thus $f_{a,j} = f^{b,j}$ on $(A' \cap \Delta_a(r_{a,j})) \times (B' \cap \Delta_b(s_{b,j})) \setminus M$. Applying Lemma 6(a), we conclude that

$$f_{a,j} = f^{b,j} \quad \text{on } (\Delta_a(r_{a,j}) \times \Delta_b(s_{b,j})) \setminus M.$$

Now let $a', a'' \in A' \cap D_j$ be such that $\Delta_{a'}(r_{a',j}) \cap \Delta_{a''}(r_{a'',j}) \neq \emptyset$. Fix a $b \in B' \cap G_j$. We know that $f_{a',j} = f^{b,j} = f_{a'',j}$ on $(\Delta_{a'}(r_{a',j}) \cap \Delta_{a''}(r_{a'',j})) \times \Delta_b(s_{b,j}) \setminus M$. Hence, by the identity principle, we conclude that $f_{a',j} = f_{a'',j}$ on $(\Delta_{a'}(r_{a',j}) \cap \Delta_{a''}(r_{a'',j})) \times G_j \setminus M$.

The same argument works for $b', b'' \in B' \cap G_j$.

Consequently, we obtain a function $f_j \in \mathcal{O}(\widetilde{U}_j \setminus M)$ with $f_j = f$ on $X'_j \setminus M$.

Let U_j be the connected component of $\widetilde{U}_j \cap \widehat{X}'_j$ with $X'_j \subset U_j$. Thus we have $f_j \in \mathcal{O}(U_j \setminus M)$ with $f_j = f$ on $X'_j \setminus M$.

Recall that $X'_j \subset U_j \subset \widehat{X}'_j$. We claim that the envelope of holomorphy of U_j coincides with $\widehat{X}'_j$. In fact, let $h \in \mathcal{O}(U_j)$, then $h|_{X'_j} \in \mathcal{O}_s(X'_j)$. So, in virtue of

Theorem 1, there exists an $\widehat{h} \in \mathcal{O}(\widehat{X}'_j)$ with $\widehat{h} = h$ on X'_j . Lemma 6(b₂) implies that $\widehat{h} = h$ on U_j .

Applying the Grauert–Remmert theorem (cf. [Jar-Pfl 2000], Th. 3.4.7), we find a function $\widehat{f}_j \in \mathcal{O}(\widehat{X}'_j \setminus M)$ with $\widehat{f}_j = f_j$ on $U_j \setminus M$. In particular, $\widehat{f}_j = f$ on $X'_j \setminus M$.

Since $A \setminus A'$, $B \setminus B'$ are pluripolar, we get

$$\begin{aligned}\widehat{X}'_j &= \{(z, w) \in D_j \times G_j : h_{A' \cap D_j, D_j}^*(z) + h_{B' \cap G_j, G_j}^*(w) < 1\} \\ &= \{(z, w) \in D_j \times G_j : h_{A \cap D_j, D_j}^*(z) + h_{B \cap G_j, G_j}^*(w) < 1\} = \widehat{X}_j.\end{aligned}$$

So, in fact, $\widehat{f}_j \in \mathcal{O}(\widehat{X}_j \setminus M)$. Using Lemma 6(b₁), we even see that $\widehat{f}_j = f$ on $X_j \setminus M$.

Observe that $\bigcup_{j=1}^{\infty} X_j = X$, $\widehat{X}_j \subset \widehat{X}_{j+1}$, and $\bigcup_{j=1}^{\infty} \widehat{X}_j = \widehat{X}$. Using again Lemma 6(a), by gluing the $\widehat{f}_j$'s, we get a function $\widehat{f} \in \mathcal{O}(\widehat{X} \setminus M)$ with $\widehat{f} = f$ on $X \setminus M$, which finishes the proof of Step 4. $\square$

To apply Step 4 we introduce the following condition (*):

Let $\rho > 0, 0 < r < R$. Put

$$\Omega := \Delta_{a_0}(\rho) \times \Delta_{b_0}(R) \subset \mathbb{C}^p \times \mathbb{C}^q, \quad \widetilde{\Omega} := \Delta_{a_0}(\rho) \times \Delta_{b_0}(r) \subset \mathbb{C}^p \times \mathbb{C}^q.$$

Let $A \subset \Delta_{a_0}(\rho) \subset \mathbb{C}^p$ be locally pluriregular, $a_0 \in A$, and let M be a pure one-codimensional analytic subset of Ω with $M \cap \widetilde{\Omega} = \emptyset$. Put $M_a := \{w \in \Delta_{b_0}(R) : (a, w) \in M\}$, $a \in A$.

Then we say that the condition (*) holds if:

For any $R' \in (r, R)$ there exists $\rho' \in (0, \rho)$ such that for any function $f \in \mathcal{O}(\widetilde{\Omega})$ with $f(a, \cdot) \in \mathcal{O}(\Delta_{b_0}(R) \setminus M_a)$, $a \in A$, there exists an extension $\widehat{f} \in \mathcal{O}(\Delta_{a_0}(\rho') \times \Delta_{b_0}(R') \setminus M)$ with $\widehat{f} = f$ on $\Delta_{a_0}(\rho') \times \Delta_{b_0}(r)$.

Step 5. *If condition (*) is satisfied, then the assumptions of Step 4 are fulfilled.*

Proof of Step 5. Take $X, M, f \in \mathcal{O}_s(X \setminus M)$ as is in Step 4. Define

$$A' := \{a \in A : M_a \neq G\}, \quad B' := \{a \in B : M^b \neq D\},$$

where $M_a := \{w \in G : (a, w) \in M\}$, $M^b := \{z \in D : (z, b) \in M\}$. It clear that $A \setminus A'$, $B \setminus B'$ are pluripolar.

Let $(D_j)_{j=1}^{\infty}, (G_j)_{j=1}^{\infty}$ be approximation sequences: $D_j \Subset D_{j+1} \Subset D$, $G_j \Subset G_{j+1} \Subset G$, $D_j \nearrow D$, $G_j \nearrow G$, $A' \cap D_j \neq \emptyset$, and $B' \cap G_j \neq \emptyset$, $j \in \mathbb{N}$.

Fix a $j \in \mathbb{N}$, $a \in A' \cap D_j$ and let Ω_j be the set of all $b \in G_{j+1}$ such that there exist a polydisc $\Delta_{(a,b)}(r_b) \subset D_j \times G_{j+1}$ and a function $\widetilde{f}_b \in \mathcal{O}(\Delta_{(a,b)}(r_b) \setminus M)$ with $\widetilde{f}_b = f$ on $(A \cap \Delta_a(r_b)) \times \Delta_b(r_b) \setminus M$.

It is clear that Ω_j is open. Observe that $\Omega_j \neq \emptyset$. Indeed, since $B \cap G_j \setminus M_a \neq \emptyset$, we find a point $b \in B \cap G_j \setminus M_a$. Therefore there is a polydisc $\Delta_{(a,b)}(r) \subset D_j \times G_j \setminus M$. Put

$$Y := \mathbb{X}(A \cap \Delta_a(r), B \cap \Delta_b(r); \Delta_a(r), \Delta_b(r)).$$

By Theorem 1, we find $r_b \in (0, r)$ and $\tilde{f}_b \in \mathcal{O}(\Delta_{(a,b)}(r_b))$ with $\tilde{f}_b = f$ on $\Delta_{(a,b)}(r_b) \cap Y \supset (A \cap \Delta_a(r_b)) \times \Delta_b(r_b)$. Consequently, $b \in \Omega_j$.

Moreover, Ω_j is relatively closed in G_{j+1} . Indeed, let c be an accumulation point of Ω_j in G_{j+1} and let $\Delta_c(3R) \subset G_{j+1}$. Take a point $b \in \Omega_j \cap \Delta_c(R) \setminus M_a$ and let $r \in (0, r_b]$, $r < 2R$, be such that $\Delta_{(a,b)}(r) \cap M = \emptyset$. Observe that $\tilde{f}_b \in \mathcal{O}(\Delta_{(a,b)}(r))$ and $\tilde{f}_b(z, \cdot) = f(z, \cdot) \in \mathcal{O}(\Delta_b(2R) \setminus M_z)$ for any $z \in A \cap \Delta_a(r)$. Hence, by (*) (with $R' := R$), there exists an extension $\hat{\tilde{f}}_b \in \mathcal{O}(\Delta_a(\rho') \times \Delta_b(R) \setminus M)$ ($\rho' \in (0, r)$) such that $\hat{\tilde{f}}_b = \tilde{f}_b$ on $\Delta_{(a,b)}(r)$. Take an $r_c > 0$ so small that $\Delta_{(a,c)}(r_c) \subset \Delta_a(\rho') \times \Delta_b(R)$ and put $\tilde{f}_c := \hat{\tilde{f}}_b$ on $\Delta_{(a,c)}(r_c) \setminus M$. Obviously $\tilde{f}_c = \hat{\tilde{f}}_b = f$ on $(A \cap \Delta_a(r_c)) \times \Delta_c(r_c) \setminus M$. Hence $c \in \Omega_j$.

Thus $\Omega_j = G_{j+1}$. There exists a finite set $T \subset \overline{G_j}$ such that

$$\overline{G_j} \subset \bigcup_{b \in T} \Delta_b(r_b).$$

Define $r_{a,j} := \min\{r_b : b \in T\}$. Take $b', b'' \in T$ with $\Delta_{b'}(r_{b'}) \cap \Delta_{b''}(r_{b''}) \neq \emptyset$. Then $\tilde{f}_{b'} = f = \tilde{f}_{b''}$ on $(A' \cap \Delta_a(r_{a,j})) \times (\Delta_{b'}(r_{b'}) \cap \Delta_{b''}(r_{b''})) \setminus M$. Consequently, by Lemma 6(a), $\tilde{f}_{b'} = \tilde{f}_{b''}$ on $\Delta_a(r_{a,j}) \times (\Delta_{b'}(r_{b'}) \cap \Delta_{b''}(r_{b''})) \setminus M$. In particular, by gluing the functions $(\tilde{f}_b)_{b \in T}$, we get a function $f_{a,j} \in \mathcal{O}(\Delta_a(r_{a,j}) \times G_j \setminus M)$ such that $f_{a,j} = f$ on $(A' \cap \Delta_a(r_{a,j})) \times G_j \setminus M$.

Changing the role of z and w , we get $f^{b,j}$, $b \in B' \cap G_j$.

Thus the assumptions of Step 4 are fulfilled. $\square$

It remains to check (*).

Step 6. *The condition (*) is always satisfied, i.e. the Main Theorem is true.*

Proof of Step 6. Fix a function $f \in \mathcal{O}(\tilde{\Omega})$ such that $f(a, \cdot) \in \mathcal{O}(\Delta_{b_0}(R) \setminus M_a)$ for any $a \in A$ with $M_a \neq \Delta_{b_0}(R)$. Define

$$R_0^* := \sup\{R' \in [r, R) : \exists_{\rho' \in (0, \rho]} \exists_{\hat{f} \in \mathcal{O}(\Delta_{a_0}(\rho') \times \Delta_{b_0}(R') \setminus M)} : \hat{f} = f \text{ on } \Delta_{a_0}(\rho') \times \Delta_{b_0}(r)\}. \quad (3)$$

It suffices to show that $R_0^* = R$.

Suppose that $R_0^* < R$. Fix $R_0^* < R'_0 < R_0 < R$ and choose $R', \rho', \hat{f}$ as in (3) with $R' \in [r, R_0^*)$, $\sqrt[q]{R'^{q-1} R'_0} > R_0^*$.

Write $w = (w', w_q) \in \mathbb{C}^q = \mathbb{C}^{q-1} \times \mathbb{C}$. Put $\tilde{A} := A \cap \Delta_{a_0}(\rho')$.

Let A' denote the set of all $(a, b') \in \tilde{A} \times \Delta_{b'_0}(R')$ which satisfy the following condition:

(*) There exist $R'' \in (R_0, R)$, $\delta > 0$, $m \in \mathbb{N}$, $c_1, \dots, c_m \in \Delta_{b_{0,q}}(R'')$, $\varepsilon > 0$, and holomorphic functions $\varphi_\mu: \Delta_{(a,b')}(\delta) \longrightarrow \Delta_{c_\mu}(\varepsilon)$, $\mu = 1, \dots, m$, such that:

- $\Delta_{(a,b')}(\delta) \subset \Delta_{a_0}(\rho') \times \Delta_{b'_0}(R')$,
- $\Delta_{c_\mu}(\varepsilon) \Subset \Delta_{b_{0,q}}(R'')$, $\mu = 1, \dots, m$,
- $\overline{\Delta_{c_\mu}}(\varepsilon) \cap \overline{\Delta_{c_\nu}}(\varepsilon) = \emptyset$ for $\mu \neq \nu$, $\mu, \nu = 1, \dots, m$,
- $\tilde{H} := \Delta_{b_{0,q}}(R') \cap H \neq \emptyset$, where $H := \Delta_{b_{0,q}}(R'') \setminus \bigcup_{\mu=1}^m \overline{\Delta_{c_\mu}}(\varepsilon)$,
- $(\Delta_{(a,b')}(\delta) \times \Delta_{b_{0,q}}(R'')) \cap M = \bigcup_{\mu=1}^m \{(z, w', \varphi_\mu(z, w')) : (z, w') \in \Delta_{(a,b')}(\delta)\}$.

For any $(a, b') \in A'$ define a new cross

$$Y := \mathbb{X}((A \cap \Delta_a(\delta)) \times \Delta_{b'}(\delta), \tilde{H}; \Delta_{(a,b')}(\delta), H).$$

Notice that Y does not intersect M . In particular, $\hat{f}|_Y \in \mathcal{O}_s(Y)$. Hence, by Theorem 1, there exists an $\hat{f}_1 \in \mathcal{O}(\hat{Y})$ with $\hat{f}_1 = \hat{f}$ on Y . Take $R''' \in (R_0, R'')$, and $\varepsilon'' > \varepsilon' > \varepsilon$ ($\varepsilon'' \approx \varepsilon$), such that

- $\Delta_{c_\mu}(\varepsilon'') \Subset \Delta_{b_{0,q}}(R''')$, $\mu = 1, \dots, m$,
- $\overline{\Delta_{c_\mu}}(\varepsilon'') \cap \overline{\Delta_{c_\nu}}(\varepsilon'') = \emptyset$ for $\mu \neq \nu$, $\mu, \nu = 1, \dots, m$.

Then there exists $\delta' \in (0, \delta]$ such that

- $\Delta_{(a,b')}(\delta') \times H' \subset \hat{Y}$, where $H' := \Delta_{b_{0,q}}(R''') \setminus \bigcup_{\mu=1}^m \overline{\Delta_{c_\mu}}(\varepsilon')$.

In particular, $\hat{f}_1 \in \mathcal{O}(\Delta_{(a,b')}(\delta') \times H')$.

Fix a $\mu \in \{1, \dots, m\}$. Then $\hat{f}_1 \in \mathcal{O}(\Delta_{(a,b')}(\delta') \times (\Delta_{c_\mu}(\varepsilon'') \setminus \overline{\Delta_{c_\mu}}(\varepsilon')))$ and $\hat{f}_1(z, w', \cdot) \in \mathcal{O}(\Delta_{c_\mu}(\varepsilon'') \setminus \{\varphi_\mu(z, w')\})$ for any $(z, w') \in (A \cap \Delta_a(\delta')) \times \Delta_{b'}(\delta')$. Using the biholomorphic mapping

$$\begin{aligned} \Phi_\mu: \Delta_{(a,b')}(\delta') \times \mathbb{C} &\longrightarrow \Delta_{(a,b')}(\delta') \times \mathbb{C}, \\ \Phi_\mu(z, w', w_q) &:= (z, w', w_q - \varphi_\mu(z, w')), \end{aligned}$$

we see that the function $g := \hat{f}_1 \circ \Phi_\mu^{-1}$ is holomorphic in $\Delta_{(a,b')}(\delta'') \times (\Delta_0(\eta'') \setminus \overline{\Delta_0}(\eta'))$ for some $\delta'' \in (0, \delta']$ and $\varepsilon' < \eta' < \eta'' < \varepsilon''$. Moreover, $g(z, w', \cdot) \in \mathcal{O}(\Delta_0(\eta'') \setminus \{0\})$ for any $(z, w') \in (A \cap \Delta_a(\delta'')) \times \Delta_{b'}(\delta'')$. Using Theorem 1 for the cross

$$\mathbb{X}((A \cap \Delta_a(\delta'')) \times \Delta_{b'}(\delta''), \Delta_0(\eta'') \setminus \overline{\Delta_0}(\eta'); \Delta_{(a,b')}(\delta''), \Delta_0(\eta'') \setminus \{0\}),$$

immediately shows that g extends holomorphically to $\Delta_{(a,b')}(\delta'') \times (\Delta_0(\eta'') \setminus \{0\})$ (because $h_{\Delta_0(\eta'') \setminus \{0\}, \Delta_0(\eta'') \setminus \overline{\Delta_0}(\eta')}^* \equiv 0$).

Transforming the above information back via Φ_μ for all μ , we conclude that the function $\hat{f}_1$ extends holomorphically to $\Delta_{(a,b')}(\delta''') \times \Delta_{b_{0,q}}(R''') \setminus M$ for some $\delta''' \in (0, \delta'']$; in particular, $\hat{f}_1$ extends holomorphically to $\Delta_{(a,b')}(\delta''') \times \Delta_{b_{0,q}}(R_0) \setminus M$.

Now we prove that $(\tilde{A} \times \Delta_{b'_0}(R')) \setminus A'$ is pluripolar. Write

$$M \cap (\Delta_{a_0}(\rho') \times \Delta_{b'_0}(R') \times \Delta_{b_{0,q}}(R)) = \bigcup_{\nu=1}^{\infty} \{\zeta \in P_\nu : g_\nu(\zeta) = 0\},$$

where $P_\nu \subseteq \Delta_{a_0}(\rho') \times \Delta_{b'_0}(R') \times \Delta_{b_{0,q}}(R)$ is a polydisc and $g_j \in \mathcal{O}(P_j)$ is a defining function for $M \cap P_j$; cf. [Chi 1989], § 2.9. Define

$$S_\nu := \{\zeta = (\tilde{\zeta}, \zeta_{p+q}) \in P_\nu : g_\nu(\zeta) = \frac{\partial g_\nu}{\partial \zeta_{p+q}}(\zeta) = 0\}$$

and observe that, by the implicit function theorem, any point from

$$(\tilde{A} \times \Delta_{b'_0}(R')) \setminus \bigcup_{\nu=1}^{\infty} \text{pr}_{\tilde{\zeta}}(S_\nu)$$

satisfies $(*)$. It is enough to show that each set $\text{pr}_{\tilde{\zeta}}(S_\nu)$ is pluripolar. Fix a ν . Let S be an irreducible component of S_ν . We have to show that $\text{pr}_{\tilde{\zeta}}(S)$ is pluripolar. If S has codimension ≥ 2 , then $\text{pr}_{\tilde{\zeta}}(S)$ is contained in a countable union of proper analytic sets (cf. [Chi 1989], § 3.8). Consequently, $\text{pr}_{\tilde{\zeta}}(S)$ is pluripolar. Thus we may assume that S is pure one-codimensional. The same argument as above shows that $\text{pr}_{\tilde{\zeta}}(\text{Sing}(S))$ is pluripolar. It remains to prove that $\text{pr}_{\tilde{\zeta}}(\text{Reg}(S))$ is pluripolar. Since g_ν is a defining function, for any $\zeta \in \text{Reg}(S)$ there exists a $k \in \{1, \dots, p+q-1\}$ such that $\frac{\partial g_\nu}{\partial \zeta_k}(\zeta) \neq 0$. Thus

$$\text{Reg}(S) = \bigcup_{k=1}^{p+q-1} T_k,$$

where $T_k := \{\zeta \in \text{Reg}(S) : \frac{\partial g_\nu}{\partial \zeta_k}(\zeta) \neq 0\}$. We only need to prove that each set $\text{pr}_{\tilde{\zeta}}(T_k)$ is pluripolar, $k = 1, \dots, p+q-1$. Fix a k . To simplify notation, assume that $k = 1$. Observe that, by the implicit function theorem, we can write

$$T_1 = \bigcup_{\ell=1}^{\infty} \{\zeta \in Q_\ell : \zeta_1 = \psi_\ell(\zeta_2, \dots, \zeta_{p+q})\},$$

where $Q_\ell \subset P_\nu$ is a polydisc, $Q_\ell = Q'_\ell \times Q''_\ell \subset \mathbb{C} \times \mathbb{C}^{p+q-1}$, and $\psi_\ell : Q''_\ell \rightarrow Q'_\ell$ is holomorphic, $\ell \in \mathbb{N}$. It suffices to prove that the projection of each set $T_{1,\ell} := \{\zeta \in Q_\ell : \zeta_1 = \psi_\ell(\zeta_2, \dots, \zeta_{p+q})\}$ is pluripolar. Fix an ℓ . Since

$$g_\nu(\psi_\ell(\zeta_2, \dots, \zeta_{p+q}), \zeta_2, \dots, \zeta_{p+q}) = 0, \quad (\zeta_2, \dots, \zeta_{p+q}) \in Q''_\ell,$$

we conclude that $\frac{\partial \psi_\ell}{\partial \zeta_{p+q}} \equiv 0$ and consequently ψ_ℓ is independent of ζ_{p+q} . Thus $\text{pr}_{\tilde{\zeta}}(T_{1,\ell}) = \{\zeta_1 = \psi_\ell(\zeta_2, \dots, \zeta_{p+q-1})\}$ and therefore the projection is pluripolar.

The proof that $(\tilde{A} \times \Delta_{b'_0}(R')) \setminus A'$ is pluripolar is completed.

Using Step 4, we conclude that $\widehat{f}$ extends holomorphically to the domain $\widehat{Y} \setminus M$, where

$$\begin{aligned}\widehat{Y} &:= \{(z, w', w_q) \in \Delta_{a_0}(\rho') \times \Delta_{b'_0}(R') \times \Delta_{b_{0,q}}(R_0) : \\ &\quad h_{A', \Delta_{a_0}(\rho') \times \Delta_{b'_0}(R')}^*(z, w') + h_{\Delta_{b_{0,q}}(R'), \Delta_{b_{0,q}}(R_0)}^*(w_q) < 1\} \\ &= \{(z, w', w_q) \in \Delta_{a_0}(\rho') \times \Delta_{b'_0}(R') \times \Delta_{b_{0,q}}(R_0) : \\ &\quad h_{\widetilde{A}, \Delta_{a_0}(\rho')}^*(z) + h_{\Delta_{b_{0,q}}(R'), \Delta_{b_{0,q}}(R_0)}^*(w_q) < 1\}\end{aligned}$$

(here we have used the product property of the relative extremal function). Since $R'_0 < R_0$, we find a $\rho_q \in (0, \rho']$ and a function $\widetilde{f}_q \in \mathcal{O}(\Delta_{a_0}(\rho_q) \times \Delta_{b'_0}(R') \times \Delta_{b_{0,q}}(R'_0) \setminus M)$ such that

$$\widetilde{f}_q = \widehat{f} \text{ on } \Delta_{a_0}(\rho_q) \times \Delta_{b_0}(R') \setminus M.$$

If $q = 1$ we get a contradiction (because $R'_0 > R_0^*$).

Let $q \geq 2$. Repeating the above argument for the coordinates w_ν , $\nu = 1, \dots, q-1$, we find a $\rho_0 \in (0, \rho']$ and a function $\widetilde{f}$ holomorphic in

$$\Delta_{a_0}(\rho_0) \times \left(\bigcup_{\nu=1}^q \Delta_{(b_{0,1}, \dots, b_{0,\nu-1})}(R') \times \Delta_{b_{0,\nu}}(R'_0) \times \Delta_{(b_{0,\nu+1}, \dots, b_{0,q})}(R') \right) \setminus M$$

such that $\widetilde{f} = \widehat{f}$ on $\Delta_{a_0}(\rho_0) \times \Delta_{b_0}(R') \setminus M$. Let $\mathcal{H}$ denote the envelope of holomorphy of the domain

$$\bigcup_{\nu=1}^q \Delta_{(b_{0,1}, \dots, b_{0,\nu-1})}(R') \times \Delta_{b_{0,\nu}}(R'_0) \times \Delta_{(b_{0,\nu+1}, \dots, b_{0,q})}(R').$$

Applying the Grauert–Remmert theorem, we can extend $\widetilde{f}$ holomorphically to $\Delta_{a_0}(\rho_0) \times \mathcal{H} \setminus M$, i.e. there exists an $\widehat{\widetilde{f}} \in \mathcal{O}(\Delta_{a_0}(\rho_0) \times \mathcal{H} \setminus M)$ with $\widehat{\widetilde{f}} = \widetilde{f}$ on $\Delta_{a_0}(\rho_0) \times \Delta_{b_0}(r)$. Observe that $\Delta_{b_0}(\sqrt[q]{R'^{q-1}R'_0}) \subset \mathcal{H}$. Recall that $\sqrt[q]{R'^{q-1}R'_0} > R_0^*$; contradiction. $\square$

Remark. Notice that the proof of Step 6 shows that the following stronger version of (*) is true: Let $\rho > 0, 0 < r < R$, $\Omega, \widetilde{\Omega}, A$, and a be as in (*). Let M be a pure one-codimensional analytic subset of Ω (we do not assume that $M \cap \widetilde{\Omega} = \emptyset$). Then:

For any $R' \in (r, R)$ there exists $\rho' \in (0, \rho)$ such that for any function $f \in \mathcal{O}(\widetilde{\Omega} \setminus M)$ with $f(a, \cdot) \in \mathcal{O}(\Delta_{b_0}(R) \setminus M_a)$, $a \in A$, there exists an extension $\widehat{f} \in \mathcal{O}(\Delta_{a_0}(\rho') \times \Delta_{b_0}(R') \setminus M)$ with $\widehat{f} = f$ on $\Delta_{a_0}(\rho') \times \Delta_{b_0}(r) \setminus M$.

4. Proof of the Main Theorem in the general case. First observe that the function $\widehat{f}$ is uniquely determined (cf. § 3).

We proceed by induction on N .

Let $D_{j,k} \nearrow D_j$, $D_{j,k} \Subset D_{j,k+1} \Subset D_j$, where $D_{j,k}$ are pseudoconvex domains with $A_{j,k} := A_j \cap D_{j,k} \neq \emptyset$. Put

$$X_k := \mathbb{X}(A_{1,k}, \dots, A_{N,k}; D_{1,k}, \dots, D_{N,k}) \subset X.$$

It suffices to show that for each $k \in \mathbb{N}$ the following condition $(**)_*$ holds.

$(**)_*$ There exists a domain U_k , $X_k \subset U_k \subset U \cap \widehat{X}_k$, such that for any $f \in \mathcal{O}_s(X \setminus M)$ there exists an $\tilde{f}_k \in \mathcal{O}(U_k \setminus M)$ with $\tilde{f}_k|_{X_k \setminus M} = f|_{X_k \setminus M}$.

Indeed, fix a $k \in \mathbb{N}$ and observe that $\widehat{X}_k$ is the envelope of holomorphy of U_k (cf. the proof of Step 4). Hence, in virtue of the Dloussky theorem (cf. [Jar-Pfl 2000], Th. 3.4.8, see also [Por 2001]), there exists an analytic subset $\widetilde{M}_k$ of $\widehat{X}_k$, $\widetilde{M}_k \cap U_k \subset M$, such that $\widehat{X}_k \setminus \widetilde{M}_k$ is the envelope of holomorphy of $U_k \setminus M$. In particular, for each $f \in \mathcal{O}_s(X \setminus M)$ there exists an $\tilde{f}_k \in \mathcal{O}(\widehat{X}_k \setminus \widetilde{M}_k)$ with $\tilde{f}_k|_{U_k \setminus M} = f|_{U_k \setminus M}$. Let $\mathcal{F}_k := \{\tilde{f}_k : f \in \mathcal{O}_s(X \setminus M)\} \subset \mathcal{O}(\widehat{X}_k \setminus \widetilde{M}_k)$. It is known that (cf. [Jar-Pfl 2000], Prop. 3.4.5) there exists a pure one-codimensional analytic subset $\widehat{M}_k \subset \widehat{X}_k$, $\widetilde{M}_k \subset \widehat{M}_k$, such that any point of $\widehat{M}_k$ is singular with respect to $\mathcal{F}_k$, i.e.

- any function $\tilde{f}_k$ extends to a function $\widehat{f}_k \in \mathcal{O}(\widehat{X}_k \setminus \widehat{M}_k)$ and
- for any $a \in \widehat{M}_k$ and an open neighborhood V of a , $V \subset \widehat{X}_k$, there exists an $f \in \mathcal{O}_s(X \setminus M)$ such that $\tilde{f}_k|_{V \setminus \widehat{M}_k}$ cannot be holomorphically extended to the whole V .

In particular, $\widehat{M}_{k+1} \cap \widehat{X}_k = \widehat{M}_k$. Consequently, $\widehat{M} := \bigcup_{k=1}^{\infty} \widehat{M}_k$ is a pure one-codimensional analytic subset of $\widehat{X}$, $\widehat{M} \cap \bigcup_{k=1}^{\infty} U_k \subset M$, and for each $f \in \mathcal{O}_s(X \setminus M)$, the function $\widehat{f} := \bigcup_{k=1}^{\infty} \widehat{f}_k$ is holomorphic on $\widehat{X} \setminus \widehat{M}$ with $\widehat{f}|_{X \setminus M} = f$.

It remains to prove $(**)_*$. Fix a $k \in \mathbb{N}$. For any $a = (a_1, \dots, a_N) \in A_{1,k} \times \dots \times A_{N,k}$ let $\rho = \rho_k(a)$ be such that $\Delta_a(\rho) \subset D_{1,k} \times \dots \times D_{N,k}$.

If $N \geq 4$, then we additionally define $(N-2)$ -fold crosses

$$Y_{k,\mu,\nu} := \mathbb{X}(A_{1,k}, \dots, A_{\mu-1,k}, A_{\mu+1,k}, \dots, A_{\nu-1,k}, A_{\nu+1,k}, \dots, A_{N,k}; \\ D_{1,k}, \dots, D_{\mu-1,k}, D_{\mu+1,k}, \dots, D_{\nu-1,k}, D_{\nu+1,k}, \dots, D_{N,k}), \\ 1 \leq \mu < \nu \leq N,$$

and we assume that ρ is so small that

$$\Delta_{(a_1, \dots, a_{\mu-1}, a_{\mu+1}, \dots, a_{\nu-1}, a_{\nu+1}, \dots, a_N)}(\rho) \subset \widehat{Y}_{k,\mu,\nu}, \quad 1 \leq \mu < \nu \leq N.$$

Since $\{(a_1, \dots, a_{j-1})\} \times \overline{D}_{j,k+1} \times \{(a_{j+1}, \dots, a_N)\} \Subset U$, we may assume that

$$\Delta_{(a_1, \dots, a_{j-1})}(\rho) \times D_{j,k+1} \times \Delta_{(a_{j+1}, \dots, a_N)}(\rho) \subset U, \quad j = 1, \dots, N. \quad (4)$$

We define N -fold crosses

$$\begin{aligned} Z_{k,a,j} &:= \mathbb{X}(A_1 \cap \Delta_{a_1}(\rho), \dots, A_{j-1} \cap \Delta_{a_{j-1}}(\rho), A_{j,k+1}, A_{j+1} \cap \Delta_{a_{j+1}}(\rho), \dots, \\ &\quad A_N \cap \Delta_{a_N}(\rho); \Delta_{a_1}(\rho), \dots, \Delta_{a_{j-1}}(\rho), D_{j,k+1}, \Delta_{a_{j+1}}(\rho), \dots, \Delta_{a_N}(\rho)), \\ &\quad j = 1, \dots, N. \end{aligned}$$

Note that $\widehat{Z}_{k,a,j} \subset U$. Since $\{(a_1, \dots, a_{j-1})\} \times \overline{D}_{j,k} \times \{(a_{j+1}, \dots, a_N)\} \Subset \widehat{Z}_{k,a,j}$, there exists an $r = r_k(a)$, $0 < r \leq \rho$, so small that

$$V_{k,a,j} := \Delta_{(a_1, \dots, a_{j-1})}(r) \times D_{j,k} \times \Delta_{(a_{j+1}, \dots, a_N)}(r) \subset \widehat{Z}_{k,a,j}, \quad j = 1, \dots, N.$$

Put

$$V_k := \bigcup_{\substack{a \in A_{1,k} \times \dots \times A_{N,k} \\ j \in \{1, \dots, N\}}} V_{k,a,j}.$$

Note that $X_k \subset V_k$. Let U_k be the connected component of $V_k \cap \widehat{X}_k$ that contains X_k .

In view of (4), the Main Theorem with $U = \widehat{X}$ (which is already proved in § 3) implies that for any $f \in \mathcal{O}_s(X \setminus M)$ there exists an extension $\widehat{f}_{k,a,j} \in \mathcal{O}(\widehat{Z}_{k,a,j} \setminus M)$ of $f|_{Z_{k,a,j} \setminus M}$. It remains to glue the functions

$$\widetilde{f}_{k,a,j} := \widehat{f}_{k,a,j}|_{V_{k,a,j} \setminus M}, \quad a \in A_{1,k} \times \dots \times A_{N,k}, \quad j = 1, \dots, N;$$

then the function $\widetilde{f}_k := \left(\bigcup_{\substack{a \in A_{1,k} \times \dots \times A_{N,k} \\ j \in \{1, \dots, N\}}} \widetilde{f}_{k,a,j} \right) \Big|_{U_k \setminus M}$ gives the required extension of $f|_{X_k \setminus M}$.

To check that the gluing process is possible, let $a, b \in A_{1,k} \times \dots \times A_{N,k}$, $i, j \in \{1, \dots, N\}$ be such that $V_{k,a,i} \cap V_{k,b,j} \neq \emptyset$. We have the following two cases:

(a) $i \neq j$: We may assume that $i = N-1$, $j = N$. Write $w = (w', w'') \in \mathbb{C}^{k_1 + \dots + k_{N-2}} \times \mathbb{C}^{k_{N-1} + k_N}$. Observe that

$$V_{k,a,N-1} \cap V_{k,b,N} = \left(\Delta_{a'}(r_k(a)) \cap \Delta_{b'}(r_k(b)) \right) \times \Delta_{b_{N-1}}(r_k(b)) \times \Delta_{a_N}(r_k(a)).$$

For $c = (c', c'')$, let

$$\begin{aligned} M_{c'} &:= \{w'' \in \mathbb{C}^{k_{N-1} + k_N} : (c', w'') \in M\}, \\ M^{c''} &:= \{w' \in \mathbb{C}^{k_1 + \dots + k_{N-2}} : (w', c'') \in M\}; \end{aligned}$$

$M_{c'}$ and $M^{c''}$ are analytic subsets of

$$\begin{aligned} U_{c'} &:= \{w'' \in \mathbb{C}^{k_{N-1} + k_N} : (c', w'') \in U\}, \\ U^{c''} &:= \{w' \in \mathbb{C}^{k_1 + \dots + k_{N-2}} : (w', c'') \in U\}, \end{aligned}$$

respectively.

We consider the following three subcases:

$N = 2$: Then $V_{k,a,1} \cap V_{k,b,2} = \Delta_{b_1}(r_k(b)) \times \Delta_{a_2}(r_k(a))$. We know that $\tilde{f}_{k,a,1} = \tilde{f}_{k,b,2}$ on the non-pluripolar set $(A_1 \cap \Delta_{b_1}(r_k(b))) \times (A_2 \cap \Delta_{a_2}(r_k(a))) \setminus M$. Hence, by the identity principle, $\tilde{f}_{k,a,1} = \tilde{f}_{k,b,2}$ on $V_{k,a,1} \cap V_{k,b,2} \setminus M$.

$N = 3$: Then $V_{k,a,2} \cap V_{k,b,3} = (\Delta_{a_1}(r_k(a)) \cap \Delta_{b_1}(r_k(b)) \times \Delta_{b_2}(r_k(b)) \times \Delta_{a_3}(r_k(a)))$. Let C'' denote the set of all points $c'' \in (A_2 \cap \Delta_{b_2}(r_k(b))) \times (A_3 \cap \Delta_{a_3}(r_k(a)))$ such that the set $M^{c''}$ has codimension ≥ 1 (i.e. for any $w' \in M^{c''}$ the codimension of $M^{c''}$ at w' is ≥ 1). Note that C'' is non-pluripolar. We have $\tilde{f}_{k,a,2}(\cdot, c'') = f(\cdot, c'') = \tilde{f}_{k,b,3}(\cdot, c'')$ on $\Delta_{a_1}(r_k(a)) \cap \Delta_{b_1}(r_k(b)) \setminus M^{c''}$.

Now, let $c' \in \Delta_{a_1}(r_k(a)) \cap \Delta_{b_1}(r_k(b))$ be such that the set $M_{c'}$ has codimension ≥ 1 . Then $\tilde{f}_{k,a,2}(c', \cdot) = \tilde{f}_{k,b,3}(c', \cdot)$ on $C'' \setminus M_{c'}$. Consequently, by the identity principle, $\tilde{f}_{k,a,2}(c', \cdot) = \tilde{f}_{k,b,3}(c', \cdot)$ on $\Delta_{b_2}(r_k(b)) \times \Delta_{a_3}(r_k(a)) \setminus M_{c'}$. Finally, $\tilde{f}_{k,a,2} = \tilde{f}_{k,b,3}$ on $V_{k,a,2} \cap V_{k,b,3} \setminus M$.

If $N \in \{2, 3\}$, then we jump directly to (b) and we conclude that the Main Theorem is true for $N \in \{2, 3\}$.

$N \geq 4$: Here is the only place where the induction over N is used. We assume that the Main Theorem is true for $N - 1 \geq 3$.

Similarly as in the case $N = 3$, let C'' denote the set of all points $c'' \in (A_{N-1} \cap \Delta_{b_{N-1}}(r_k(b))) \times (A_N \cap \Delta_{a_N}(r_k(a)))$ such that the set $M^{c''}$ has codimension ≥ 1 ; C'' is non-pluripolar. The function $f_{c''} := f(\cdot, c'')$ is separately holomorphic on $Y_{k,N-1,N} \setminus M^{c''}$. By the inductive assumption, the function $f_{c''}$ extends to a function $\hat{f}_{c''} \in \mathcal{O}(\hat{Y}_{k,N-1,N} \setminus \hat{M}(c''))$, where $\hat{M}(c'')$ is an analytic subset of $\hat{Y}_{k,N-1,N}$ with $\hat{M}(c'') \subset M^{c''}$ in an open neighborhood of $Y_{k,N-1,N}$. Recall that

$$\Delta_{a'}(r_k(a)) \cup \Delta_{b'}(r_k(b)) \subset \hat{Y}_{k,N-1,N}.$$

Since $\tilde{f}_{k,a,N-1}(\cdot, c'') = f_{c''}$ on $\Delta_{a'}(r_k(a)) \cap Y_{k,N-1,N} \setminus M^{c''}$ and $\tilde{f}_{k,b,N}(\cdot, c'') = f_{c''}$ on $\Delta_{b'}(r_k(b)) \cap Y_{k,N-1,N} \setminus M^{c''}$, we conclude that $\tilde{f}_{k,a,N-1}(\cdot, c'') = \hat{f}_{c''} = \tilde{f}_{k,b,N}(\cdot, c'')$ on $\Delta_{a'}(r_k(a)) \cap \Delta_{b'}(r_k(b)) \setminus M^{c''}$.

Let $c' \in \Delta_{a'}(r_k(a)) \cap \Delta_{b'}(r_k(b))$ be such that the set $M_{c'}$ has codimension ≥ 1 . Then $\tilde{f}_{k,a,N-1}(c', \cdot) = \tilde{f}_{k,b,N}(c', \cdot)$ on $C'' \setminus M_{c'}$. Consequently, by the identity principle, $\tilde{f}_{k,a,N-1}(c', \cdot) = \tilde{f}_{k,b,N}(c', \cdot)$ on $\Delta_{b_{N-1}}(r_k(b)) \times \Delta_{a_N}(r_k(a)) \setminus M_{c'}$ and, finally, $\tilde{f}_{k,a,N-1} = \tilde{f}_{k,b,N}$ on $V_{k,a,N-1} \cap V_{k,b,N} \setminus M$.

(b) $i = j$: We may assume that $i = j = N$. Observe that

$$V_{k,a,N} \cap V_{k,b,N} = \left(\Delta_{(a_1, \dots, a_{N-1})}(r_k(a)) \cap \Delta_{(b_1, \dots, b_{N-1})}(r_k(b)) \right) \times D_{N,k}.$$

By (a) we know that:

$$\begin{aligned} \tilde{f}_{k,a,N} &= \tilde{f}_{k,a,N-1} \text{ on } V_{k,a,N} \cap V_{k,a,N-1} \setminus M, \\ \tilde{f}_{k,a,N-1} &= \tilde{f}_{k,b,N} \text{ on } V_{k,a,N-1} \cap V_{k,b,N} \setminus M. \end{aligned}$$

Hence

$$\begin{aligned}\tilde{f}_{k,a,N} &= \tilde{f}_{k,b,N} \text{ on } V_{k,a,N} \cap V_{k,a,N-1} \cap V_{k,b,N} \setminus M \\ &= \left(\Delta_{(a_1, \dots, a_{N-1})}(r_k(a)) \cap \Delta_{(b_1, \dots, b_{N-1})}(r_k(b)) \right) \times \Delta_{a_N}(r_k(a)) \setminus M,\end{aligned}$$

and finally, by the identity principle,

$$\tilde{f}_{k,a,N} = \tilde{f}_{k,b,N} \text{ on } V_{k,a,N} \cap V_{k,b,N} \setminus M.$$

The proof of the Main Theorem is completed.

5. Proof of Theorem 2. To prove Theorem 2 we need the following version of the Hartogs theorem (cf. [Chi-Sad 1988], Th. 1). Let E denote the unit disc.

Theorem 7. *Let $A \subset E^{N-1}$ be locally pluriregular, let $U \subset E^{N-1} \times \mathbb{C}$ be an open neighborhood of $A \times \mathbb{C}$, and let $M \subset U$ be a relatively closed set such that $M \cap E^N = \emptyset$ and for any $a \in A$ the fiber $M_a := \{w \in \mathbb{C} : (a, w) \in M\}$ is polar. Put $X := \mathbb{X}(A, E; E^{N-1}, \mathbb{C})$. Then there exists a relatively closed pluripolar set $S \subset E^{N-1} \times \mathbb{C}$, $S \cap E^N = \emptyset$, such that $S_a \subset M_a$, $a \in A$, and any function $f \in \mathcal{O}_s(X \setminus M)$ extends holomorphically to $\hat{X} \setminus S = E^{N-1} \times \mathbb{C} \setminus S$.*

Proof. Let $\mathcal{F} := \mathcal{O}_s(X \setminus M)$. It is known that each function $f \in \mathcal{F}$ has the univalent domain of existence $G_f \subset E^{N-1} \times \mathbb{C}$ ^(‡). Let G denote the connected component of $\text{int} \bigcap_{f \in \mathcal{F}} G_f$ that contains E^N and let $S := E^{N-1} \times \mathbb{C} \setminus G$. It remains to show that S is pluripolar.

Take $a \in A$ and $b \in \mathbb{C} \setminus M_a$. Since M_a is polar, there exists a curve $\gamma : [0, 1] \rightarrow \mathbb{C} \setminus M_a$ such that $\gamma(0) = 0$, $\gamma(1) = b$. Take an $\varepsilon > 0$ so small that

$$\Delta_a(\varepsilon) \times (\gamma([0, 1]) + \Delta_b(\varepsilon)) \subset U \setminus M.$$

Put $V_b := E \cup (\gamma([0, 1]) + \Delta_b(\varepsilon))$ and consider the cross

$$Y := \mathbb{X}(A \cap \Delta_a(\varepsilon), E; \Delta_a(\varepsilon), V_b).$$

Then $f \in \mathcal{O}_s(Y)$ for any $f \in \mathcal{F}$. Consequently, by Theorem 1, $\hat{Y} \subset G_f$ for any $f \in \mathcal{F}$ and hence $\hat{Y} \subset G$. In particular, $\{a\} \times (\mathbb{C} \setminus M_a) \subset G$.

Thus $S_a \subset M_a$ for all $a \in A$. Consequently, by Lemma 5 from [Chi-Sad 1988], S is pluripolar. $\square$

Proof of Theorem 2. Fix a point $c = (a, b) \in A \times B \setminus M$ and let $\rho > 0$ be such that $\Delta_c(\rho) \cap M = \emptyset$. Consider the cross

$$Y := \mathbb{X}(A \cap \Delta_a(\rho), B \cap \Delta_b(\rho); \Delta_a(\rho), \Delta_b(\rho))$$

(‡) We like to thank Professor Evgeni Chirka for explaining us some details of the proof of Theorem 1 in [Chi-Sad 1988].

and let $0 < r \leq \rho$ be such that $\Delta_c(r) \subset \widehat{Y}$. Put $A_0 := A \cap \Delta_a(r)$, $B_0 := B \cap \Delta_b(r)$. By Theorem 1, for any $f \in \mathcal{O}_s(X \setminus M)$ there exists an $\tilde{f} \in \mathcal{O}(\Delta_c(r))$ with $\tilde{f} = f$ on $A_0 \times B_0$.

Define

$$U := \mathbb{C} \times \Delta_b(r), \quad V := \Delta_a(r) \times \mathbb{C}.$$

Observe that the sets

$$P := \{z \in \mathbb{C} : M_z \text{ is not polar}\}, \quad Q := \{w \in \mathbb{C} : M^w \text{ is not polar}\}$$

are polar. Indeed, let $v \in \mathcal{PSH}(\mathbb{C}^2)$, $v \not\equiv -\infty$, be such that $M \subset v^{-1}(-\infty)$. Define

$$u(z) := \sup\{v(z, w) : w \in \overline{E}\}, \quad z \in \mathbb{C}.$$

Then $P \subset u^{-1}(-\infty)$. It remains to observe that $u \in \mathcal{PSH}(\mathbb{C})$ and $u \not\equiv -\infty$.

Now, by Theorem 7, there exist relatively closed pluripolar sets $S \subset U \setminus \Delta_c(r)$, $T \subset V \setminus \Delta_c(r)$, such that for any $f \in \mathcal{O}_s(X \setminus M)$ there exist $\tilde{f}_1 \in \mathcal{O}(U \setminus S)$, $\tilde{f}_2 \in \mathcal{O}(V \setminus T)$ with $\tilde{f}_1 = \tilde{f}_2 = \tilde{f}$ on $\Delta_c(r)$. Consequently, the function $\tilde{f} := \tilde{f}_1 \cup \tilde{f}_2$ is well defined on $U \cup V \setminus (S \cup T)$. Note that $R := S \cup T$ is a relatively closed pluripolar subset of $W := U \cup V$. Moreover, $\mathbb{C}^2$ is the envelope of holomorphy of W .

Now, by [Chi 1993], there exists a closed pluripolar set $\widehat{M} \subset \mathbb{C}^2$ with $\widehat{M} \cap W \subset R$ such that $\mathbb{C}^2 \setminus \widehat{M}$ is the envelope of holomorphy of $W \setminus R$. Consequently, for each $f \in \mathcal{O}_s(X \setminus M)$ there exists an $\widehat{f} \in \mathcal{O}(\mathbb{C}^N \setminus \widehat{M})$ with $\widehat{f} = f$ on $A_0 \times B_0$.

Define

$$P' := \{z \in \mathbb{C} : (M \cup \widehat{M})_z \text{ is not polar}\}, \quad Q' := \{w \in \mathbb{C} : (M \cup \widehat{M})^w \text{ is not polar}\},$$

$A' := A \setminus P'$, $B' := B \setminus Q'$, $X' := \mathbb{X}(A', B'; \mathbb{C}, \mathbb{C})$. Now we argue as in the proof of Theorem 1.1 from [Sic 2000] and we conclude that $\widehat{f} = f$ on $X' \setminus (M \cup \widehat{M})$. $\square$

REFERENCES

- [Ale-Zer 2001] O. Alehyane, A. Zeriahi, *Une nouvelle version du théorème d'extension de Hartogs pour les applications séparément holomorphes entre espaces analytiques*, Ann. Polon. Math. 76 (2001), 245–278.
- [Chi 1989] E. M. Chirka, *Complex Analytic Sets*, Kluwer Acad. Publishers, 1989.
- [Chi 1993] E. M. Chirka, *The extension of pluripolar singularity sets*, Proc. Steklov Inst. Math. 200 (1993), 369–373.
- [Chi-Sad 1988] E. M. Chirka, A. Sadullaev, *On continuation of functions with polar singularities*, Math. USSR-Sb. 60 (1988), 377–384.
- [Jar-Pfl 2000] M. Jarnicki, P. Pflug, *Extension of Holomorphic Functions*, de Gruyter Expositions in Mathematics 34, Walter de Gruyter, 2000.
- [Jar-Pfl 2001] M. Jarnicki, P. Pflug, *Cross theorem*, IMUJ Preprint 2001/08 (to appear in Ann. Polon. Math. (2001)).
- [Kli 1991] M. Klimek, *Pluripotential Theory*, Oxford University Press, 1991.
- [Ngu 1997] Nguyen Thanh Van, *Separate analyticity and related subjects*, Vietnam J. Math. 25 (1997), 81–90.

- [Ngu-Sic 1991] Nguyen Thanh Van, J. Siciak Fonctions plurisousharmoniques extrémales et systèmes doublement orthogonaux de fonctions analytiques, *Bull. Sc. Math.* 115 (1991), 235–244.
- [Ngu-Zer 1991] Nguyen Thanh Van, A. Zeriahi, *Une extension du théorème de Hartogs sur les fonctions séparément analytiques*, *Analyse Complexe Multivariables, Récents Développements*, A. Meril (ed.), EditEl, Rende, 1991, pp. 183–194.
- [Ngu-Zer 1995] Nguyen Thanh Van, A. Zeriahi, *Systèmes doublement orthogonaux de fonctions holomorphes et applications*, *Banach Center Publ.* 31 (1995), 281–297.
- [Ökt 1998] O. Öktem, *Extension of separately analytic functions and applications to range characterization of exponential Radon transform*, *Ann. Polon. Math.* 70 (1998), 195–213.
- [Ökt 1999] O. Öktem, *Extending separately analytic functions in $\mathbb{C}^{n+m}$ with singularities*, *Extension of separately analytic functions and applications to mathematical tomography (Thesis)*, Dep. Math. Stockholm Univ., 1999.
- [Por 2001] E. Porten, *On the Hartogs–phenomenon and extension of analytic hypersurfaces in non-separated Riemann domains*, Preprint (2001).
- [Shi 1989] B. Shiffman, *On separate analyticity and Hartogs theorem*, *Indiana Univ. Math. J.* 38 (1989), 943–957.
- [Sic 1969] J. Siciak, *Separately analytic functions and envelopes of holomorphy of some lower dimensional subsets of $\mathbb{C}^n$* , *Ann. Polon. Math.* 22 (1969–1970), 147–171.
- [Sic 1981] J. Siciak, *Extremal plurisubharmonic functions in $\mathbb{C}^N$* , *Ann. Polon. Math.* 39 (1981), 175–211.
- [Sic 2000] J. Siciak, *Holomorphic functions with singularities on algebraic sets*, IMUJ Preprint 2000/21 (to appear in *Univ. Iag. Acta Math.* (2001)).
- [Zah 1976] V. P. Zahariuta, *Separately analytic functions, generalizations of Hartogs theorem, and envelopes of holomorphy*, *Math. USSR-Sb.* 30 (1976), 51–67.

UNIwersytet Jagielloński
 Instytut Matematyki
 30-059 Kraków, Reymonta 4, Poland
E-mail: jarnicki@im.uj.edu.pl

Carl von Ossietzky Universität Oldenburg
 Fachbereich Mathematik
 Postfach 2503
 D-26111 Oldenburg, Germany
E-mail: pflug@mathematik.uni-oldenburg.de